

Combined use of xenogenous bone blocks and guided bone regeneration for three-dimensional augmentation of anterior maxillary ridge: A case series

Mohammad Reza Talebi¹ • Noushin Janbakhsh^{2*}

¹Department of Periodontology, School of Dentistry, Shahid Beheshti University of Medical Sciences, Tehran, Iran

²Department of Periodontology, School of Dentistry, Ilam University of Medical Sciences, Ilam, Iran

*Corresponding Author; E-mail: noushinjn@yahoo.com

Received: 13 October 2019; Accepted: 30 December 2019

J Adv Periodontal Implant Dent 2019;11(2):94-98 | [doi:10.15171/japid.2019.015](https://doi.org/10.15171/japid.2019.015)

This article is available from <https://japid.tbzmed.ac.ir/>

© 2019 The Authors. This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Abstract

Background. Bone augmentation ensures a favorable 3-dimensional position of implants. Onlay grafting is one of the techniques in ridge augmentation, which can be performed with the use of xenogenous blocks.

Methods. Three cases of the vertical and horizontal ridge are discussed, which were augmented using xenogenous blocks. The blocks were shaped in a favorable size and puzzled along the grafting area. All the gaps were filled with granular xenografts. The flaps were coronally advanced to obtain primary closure.

Results. An average of 4.2-mm gain in width and 4.2-mm gain in height of the ridge was observed at the implantation stage.

Conclusion. The outcomes of these cases could pave the way for suggesting xenograft blocks for augmenting wide areas of the alveolar ridge on average of 4 mm in width and height in selected cases as an alternative to standard autogenous blocks. Long-lasting xenograft ensures implant and lip support in the esthetic zone.

Key words: Alveolar bone grafting, alveolar bone loss, Heterograft.

Introduction

Several materials and techniques have been developed to augment the alveolar bone.¹ Autogenous bone as the gold standard graft material has several disadvantages, including morbidity of the donor site, patient discomfort, unpredictable resorption of the graft, and the limited quantity.² The use of xenografts has been advocated due to the lack of the shortcomings mentioned above.³

Various techniques of ridge augmentation include guided bone regeneration, onlay/veneer grafts, inlay

grafts, distraction osteogenesis, and ridge splitting.^{1,4} Onlay xenografts are reported to result in 97.1% implant survival.⁴ In this technical note, we present three cases of vertical and horizontal ridge augmentation using xenogenous bone blocks.

Methods

Three consecutive generally healthy patients in a private periodontal office with deficient maxillary alveolar ridge <3 mm in thickness buccolingually were included (Figures 1–2). The primary defect size and location were assessed using CBCT images (Table 1).

Table 1. The overview of surgical sites augmented by onlay xenografts

Patient	Age	M/F	Ridge augmentation (xenogenous onlay grafts)			
			Graft material	Membrane type	Grafted area*	Healing (months)
1	39	F	Ceraboneblock-L20†	SIC b-mem (30*40 cm)	4-16	9
2	51	F	Ceraboneblock-L20	SIC b-mem (20*30 cm)	9-13	9
3	50	F	Ceraboneblock-L20	Jason Membrane‡ (30*40 cm)	6-14	9

*International Dental Federation tooth-numbering system.

† Healing time before uncover.

‡ Cerabone block-L20 (20*20*10mm) (aap Biomaterials)

‡ Jason Membrane (30*40cm) (aap Biomaterials)

The patients were pre-medicated with one gr of amoxicillin one hour in advance. A full-thickness flap was elevated by two incisions under local anesthesia: first, an incision on the alveolar crest, which was delicately palatal/lingual; second, a vertical releasing incision on the second tooth away from the surgical site. An incision was also made through the periosteum with a scalpel blade and Medzenbach scissors by continuously opening the scissors and cutting through the tissue attachments partially to allow coronal advancement. The advancement continued until a 2-mm overlap of the buccal flap on the palatal side was observed. Using fine tissue forceps to stretch the buccal flap to overlap the palatal side, the buccal flap should have stayed over the palatal flap to indicate that it was tension-free. However, if the buccal flap started to retract

to the buccal side, periosteal releasing was necessary to proceed.

After granulation tissue removal, the recipient site was decorticated thoroughly by a #2 or #4 round bur with a 2-mm distance between the perforations. The resorbable membrane (Table 1) was fixed labially. A xenogenous bone block (Table 1) was molded in pieces measuring about 10 mm in length, 4 mm in width, and 3-4 mm in depth, using the saw tip of a piezosurgery tool. All the pieces were adapted along the grafting area on the buccal side, and each was fixed with one or two screws (8 or 10 mm long). All the gaps between the block pieces were filled using xenogenic particulate bone graft (Cerabone, Bottis, Germany) until slightly over-contoured. After fixing another membrane palatally/lingually, the flap was




Figure 1. Case 1: 39-year-old female. a) The initial CBCT at the augmentation phase showing very poor dimensions of the bone (due to the previous traumatic extractions). b) The ridge at the augmentation session. c) Residual ridge following full-thickness flap elevation. d) CBCT of the patient 9 months later shows block integration. e) Clinical view at the implantation session, adequate height, and width of the bone and complete integration of the graft. f) Full-thickness flap. g) While preparing the implant site a trephine biopsy was obtained from the site #16. The biopsy specimen stained by hematoxylin-and-eosin and magnified $\times 200$ shows newly formed bone at the site of implant placement and remnants of the xenograft. Although more remnants and connective tissue might be present at the buccal side, it is of importance that in the favorable site of implant placement only native bone is present.

sutured first by horizontal mattress sutures and then by interrupted sutures in between. It is suggested that this suture be placed in the mucogingival area to help approximate the flaps.

The regimen of 0.2% chlorhexidine mouthwash twice a day was administered for one week, accompanied by amoxicillin (500 mg) three times a day. Ibuprofen (400 mg) was prescribed every 6 hours until the pain was relieved.

The patients were examined the day after surgery, and every 48 hours, to check and render professional cleaning. No complications, including dehiscence and infection of the surgical site, occurred during the follow-ups.

Interrupted sutures were removed by the second week, but the horizontal mattresses remained for one more week. The follow-ups continued monthly. After a healing period of 9 months, a second CBCT was taken to determine the implant size for the second-stage surgery and measurement of new bone formation.

After raising a full-thickness flap, the implants were inserted at an insertion torque of 20 N.cm and submerged for three months until they were uncovered for the prosthetic stage (Table 2). At the time of implantation, bone biopsies were gathered using a trephine bur and assessed histologically to determine graft integration.

Results

Table 3 shows the summary of the outcomes at nine months. The new bone appeared well integrated to the recipient site on CBCT images. The results of histological evaluations showed that the xenografts were integrated into the newly formed bone (Figure 1, F–G). No bone loss, peri-implant mucositis, or implant mobility were recorded.

Discussion

Xenogenous bone blocks were used to augment extensive horizontal and vertical alveolar ridge defects while managing two of the most common complications associated with them. First, the vertical releasing and periosteal incisions were made to attain tension-free primary closure. Second, piezo-surgery was used to shape the xenograft, which prevented the fracture of the fragile material.

CBCT examinations showed adequate bone gain (average: 4.4 mm horizontally and 4.2 mm vertically). Histologic evaluations showed newly formed bone at the implant site after nine months. Sufficient bone was present at the implantation site to place the implants

at the ideal site. The results reported by the sixth European Workshop on Periodontology declared 4.2–4.6 mm of increase in the vertical height of ridge after autogenous onlay grafting.⁵ Previous studies have reported different amounts of augmentation and xenograft integration in humans, as summarized in Table 4.^{6–12}

In this study, growth factors were not used; thus, the use of double-layered resorbable membranes provided a barrier during graft remodeling. This, of course, required a longer period of healing.¹³ Patients




Figure 2. Case 2 was a 51-year-old female. a) Less traumatic extraction of hopeless teeth. b) Block fixation using fixation screws; depending on the size of the blocks used one screw could provide enough fixation. c) Particulate xenograft fills the space between the block grafts. d) A double-layered membrane is fixed. e) At the implantation session, a full-thickness flap was raised, revealing complete integration of the grafts and adequate dimensions of the ridge for implant placement.

Table 2. The overview of implant sites and characteristics

Number of implants	Site*	Type	Length	Diameter
6	4,6,8 9,11,13	Tixos-MC	10 11.5	4.5
4	6, 7, 10, 11	zimmer	10	4.1
5	7, 10, 11, 12, 14	Tixos-MC	10	4.5 3.75

*International Dental Federation tooth-numbering system.

Table 3. The overview of the results: bone gain (millimeters) from CBCT tomography at the time of implantation

Patient	Last follow up (months)*	Average Width		Average Height	
		Before augmentation	After augmentation	Before augmentation	After augmentation
1	12	2.3	6.3	9	13.9
2	24	1.5	6.1	6.7	10.4
3	6	6.5	10.9	1.5	5.5
Mean	14	3.4	7.8	5.7	9.9

*The last visit after implantation by months.

also prefer bone substitutes rather than autogenous bone.¹

The PASS principles, described by Wang and Boyapati¹⁴ for GBR, can be modified by adding the factor of ‘time’ and be applied for xenogenic block grafting. The primary closure should be obtained through releasing incisions (both vertical and periosteal) and mattress and interrupted sutures. Angiogenesis is obtained by decortication of the bone. Stability would be possible by using fixation screws for blocks. The blocks themselves will provide a tenting effect for the GBR arrears,¹⁵ and finally, double-layering and screw-fixing of the membrane will provide the complex with stability. Space maintenance is achieved by the use of blocks and particulate bone grafts. Finally comes the time factor, which we believe is the key to

the success of the use of xenogenic blocks. A minimum of 9 months is required for the integration and tissue maturation in the augmented site.

Although favorable results were obtained here, augmentation procedures have high morbidity and are very skill-sensitive techniques. Clinical trials with large sample sizes are needed to confirm the results of this study.

Conclusion

Although autogenous blocks remain the standard, wide areas of bone augmentation were achieved in these cases. Primary closure, angiogenesis, stability, space maintenance, and increased healing time are the keys to successful management. The primary limitation is technique sensitivity.

Table 4. A summary of some human studies using xenograft for alveolar bone augmentation

Authors	No. of patients	Graft healing (months)	Augmentation material	Horizontal bone gain	Vertical bone gain	Graft integration
Simion et al. ⁶	7	3.5	Autog P+ Xeno P+ Ti reinforced e-PTFE mem.	-	3.15 mm	8.63% remaining xeno
Scarano et al. ⁷	9	4	Xeno miniblocks + CCPB particles	-	7.43 – 6.68 mm	33% remaining graft
Simion et al. ⁸	2	5	Xeno B/P+ rh-PDGF +/- collagen mem.	-	3 mm (with mem) 8 mm (without mem)	Xeno embedded in bone
Proussaefs and Lozada ⁹	12	5	Autog B+ Xeno P	-	5.8 mm	23.89% remaining Xeno
Friedmann et al. ¹⁰	28	7	Xeno P+ reorbable or non-resorbable mem	-	-	14 – 15% remaining Xeno
Von Arx and Buser ¹¹	42	5.8	Autog B+ Xeno P+ collagen mem	4.6 mm		Xeno particles showed either fibrous encapsulation or new bone integration Xeno integrated into new bone but on the surface of the new bone, only some single xeno particles were integrated.
Hammerle et al. ¹²	12	9.5	Xeno B/P	3.6 mm		

Xeno: Xenograft; Autog: Autogenous; P: particles; B: block; Ti: Titanium; e-PTFE: expanded-polytetrafluoroethylene; mem: membrane; CCPB: cortico-cancellous porcine bone;

+/-: with or without

Competing Interests

The authors declare no conflict(s) of interest related to the publication of this work.

Authors' Contributions

Clinical work, follow ups, design of article and intellect.

Acknowledgments

None.

Ethics Approval

None.

Funding

No funding.

References

1. Esposito M, Grusovin MG, Felice P, Karatzopoulos G, Worthington HV, Coulthard P. Interventions for replacing missing teeth: horizontal and vertical bone augmentation techniques for dental implant treatment. *Cochrane Database Syst Rev* 2009;Cd003607.
2. Khojasteh A, Kheiri L, Motamedian S, Khoshkam V. Guided bone regeneration for the reconstruction of alveolar bone defects. *Ann Maxillofac Surg* 2017; 7:263-77.
3. Simion M, Rocchietta I, Kim D, Nevins M, Fiorellini J. Vertical ridge augmentation by means of deproteinized bovine bone block and recombinant human platelet-derived growth factor-BB: a histologic study in a dog model. *Int J Periodontics Restorative Dent* 2006; 26:415-23.
4. Aghaloo TL, Moy PK. Which hard tissue augmentation techniques are the most successful in furnishing bony support for implant placement? *Int J Oral Maxillofac Implants* 2007; 22 Suppl: 49-70.
5. Tonetti MS, Hammerle CH. Advances in bone augmentation to enable dental implant placement: Consensus Report of the Sixth European Workshop on Periodontology. *J Clin Periodontol* 2008; 35:168-72.
6. Simion M, Fontana F, Rasperini G, Maiorana C. Vertical ridge augmentation by expanded-polytetrafluoroethylene membrane and a combination of intraoral autogenous bone graft and deproteinized anorganic bovine bone (Bio Oss). *Clin Oral Implants Res* 2007; 18:620-9.
7. Scarano A, Carinci F, Assenza B, Piattelli M, Murmura G, Piattelli A. Vertical ridge augmentation of atrophic posterior mandible using an inlay technique with a xenograft without miniscrews and miniplates: case series. *Clin Oral Implants Res* 2011; 22:1125-30.
8. Simion M, Rocchietta I, Dellavia C. Three-dimensional ridge augmentation with xenograft and recombinant human platelet-derived growth factor-BB in humans: report of two cases. *Int J Periodontics Restorative Dent* 2007; 27:109-15.
9. Proussaefs P, Lozada J. The use of intraorally harvested autogenous block grafts for vertical alveolar ridge augmentation: a human study. *Int J Periodontics Restorative Dent* 2005; 25:351-63.
10. Friedmann A, Strietzel FP, Marezki B, Pitaru S, Bernimoulin JP. Histological assessment of augmented jaw bone utilizing a new collagen barrier membrane compared to a standard barrier membrane to protect a granular bone substitute material. *Clin Oral Implants Res* 2002; 13:587-94.
11. von Arx T, Buser D. Horizontal ridge augmentation using autogenous block grafts and the guided bone regeneration technique with collagen membranes: a clinical study with 42 patients. *Clin Oral Implants Res* 2006; 17:359-66.
12. Hammerle CH, Jung RE, Yaman D, Lang NP. Ridge augmentation by applying bioresorbable membranes and deproteinized bovine bone mineral: a report of twelve consecutive cases. *Clin Oral Implants Res* 2008; 19:19-25.
13. Kim SH, Kim DY, Kim KH, Ku Y, Rhyu IC, Lee YM. The efficacy of a double-layer collagen membrane technique for overlaying block grafts in a rabbit calvarium model. *Clin Oral Implants Res* 2009; 20:1124-32.
14. Wang HL, Boyapati L. "PASS" principles for predictable bone regeneration. *Implant Dent* 2006; 15:8-17.
15. Khojasteh A, Hassani A, Motamedian SR, Saadat S, Alikhasi M. Cortical Bone Augmentation Versus Nerve Lateralization for Treatment of Atrophic Posterior Mandible: A Retrospective Study and Review of Literature. *Clin Implant Dent Relat Res* 2016; 18:342-59.