

Research Article



Effect of photobiomodulation on wound healing and pain after crown lengthening surgery: A randomized clinical trial

Parichehr Behfarnia¹, Reza Birang¹, Amineh Ghaznavi², Seyed Amir Mirghaderi³

¹Department of Periodontics, Dental Implant Research Center, Dental Research Institute, School of Dentistry, Isfahan University of Medical Sciences, Isfahan, Iran

²Department of Periodontics, School of Dentistry, Arak University of Medical Sciences, Arak, Iran

³School of Dentistry, Tehran University of Medical Sciences, Tehran, Iran

ARTICLE INFO

Article History:

Received: August 28, 2024

Accepted: July 9, 2025

ePublished: July 22, 2025

Keywords:

Crown lengthening, Diode laser,
Pain, Photobiomodulation,
Wound healing

Abstract

Background. The present study evaluated the synergistic effect of 808 nm and 660 nm diode lasers on the processes of healing and pain management following crown lengthening surgery.

Methods. This randomized clinical trial involved 20 patients who underwent surgical crown lengthening on both sides of their jaw. Following the surgery, one tooth from each patient was randomly assigned to either the case group (irradiated with 808 nm and 660 nm diode lasers as photobiomodulation [PBM] therapy) or the control group (laser device remained switched off). The early healing index (EHI), comprising clinical signs of inflammation (CSI), clinical signs of homeostasis (CSH), and clinical signs of re-epithelialization (CSR), was assessed on days 3 and 7. Pain severity was quantified on the day of surgery and 1, 3, and 7 days after surgery using a visual analog scale. The data were analyzed using the Wilcoxon test.

Results. No significant differences were observed in CSR on days 3 ($P=0.18$) and 7 ($P=1.0$), nor in CSI on day 3 ($P=0.477$) after surgery. However, a significant difference was identified in CSI on day 7 and in CSH on both days 3 and 7 ($P<0.05$) after surgery. Furthermore, the level of postoperative pain demonstrated a significant difference ($P\leq 0.005$).

Conclusion. PBM demonstrably enhanced CSI by day 7 and improved CSH by days 3 and 7, in addition to decreasing postoperative pain.

Introduction

Postoperative complications affect up to 15% of patients undergoing periodontal and implant surgeries.¹ The most common complications include dentinal hypersensitivity, excessive pain, postoperative bleeding, edema, and delayed wound healing.² To mitigate these issues, various approaches have been explored, with photobiomodulation (PBM), also known as low-level laser therapy (LLLT), being employed in recent years.^{3,4}

PBM elicits photochemical, photophysical, and photobiological effects within cells and tissues. The therapeutic benefits of PBM stem from its capacity for biostimulation and biomodulation at the cellular level.³ This technique employs a concentrated, low-power light beam, typically within the 600 to 1000 nm wavelength range, to facilitate tissue healing, periodontal regeneration, and anti-inflammatory responses.⁵ PBM exerts its effects by stimulating calcium channels within cell membranes and mitochondrial membrane surface receptors, thereby enhancing adenosine triphosphate (ATP) production and tissue oxygenation. Furthermore, PBM modulates reactive oxygen species, cytokine levels,

growth factors, and inflammatory mediators.^{2,5,6} The application of low-power lasers stimulates fibroblasts, keratinocytes, and collagen synthesis, angiogenesis, and enhances growth factor release, all of which collectively contribute to accelerated wound healing.⁷

Limited randomized clinical trials (RCTs) have investigated the efficacy of PBM in promoting wound healing and alleviating pain after oral surgical procedures, specifically crown lengthening. However, the findings of these RCTs have been inconsistent. Some studies have indicated that applying PBM after oral surgery culminated in improved clinical gingival healing and enhanced periodontal parameters.^{3,8} Conversely, other studies have reported no significant benefits of PBM after oral surgery for either wound healing or pain management.⁹ Furthermore, the evidence supporting the use of single or combined wavelength photons in treating periodontitis is restricted, necessitating further clinical investigations.³ Thus, the current study aimed to assess the short-term efficacy of combined 808 nm and 660 nm diode lasers in promoting healing and reducing postoperative pain after crown lengthening surgery.

*Corresponding author: Reza Birang, Email: birang@dnt.mui.ac.ir

© 2025 The Author(s). This is an open access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0/>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Methods

This research was carried out at the Department of Periodontics, Isfahan University of Medical Sciences. The research protocol received approval from the Ethics Committee of Isfahan University of Medical Sciences (IR.MUI.RESEARCH.REC.1400.128) and was subsequently registered with the Iranian Registry of Clinical Trials (IRCT20110109005570N12). All participating patients provided written informed consent, and their involvement in the study was voluntary.

Trial design

The present research was a single-center, placebo-controlled, prospective RCT, using a split-mouth approach. Figure 1 illustrates the complete study workflow. Outcome reporting adhered to the guidelines established by the Consolidated Standards of Reporting Trials (CONSORT) 2010.¹⁰

Participants, eligibility criteria, and settings

Participants were selected based on the following criteria: (I) generally healthy individuals, (II) aged 25–60 years, (III) requiring surgical crown lengthening on two non-adjacent, single- or double-rooted teeth located in

different quadrants, (IV) requiring a minimum of 2 mm of bone resection, and (V) exhibiting at least 2 mm of keratinized gingiva postoperatively.

The exclusion criteria included: (I) inadequate oral hygiene, (II) lack of cooperation or refusal to participate, (III) smoking, (IV) general contraindications for laser therapy, and (V) the presence of infection at the surgical site.

The study's sample comprised 20 consecutive patients, each undergoing crown lengthening surgery on two teeth, resulting in a total of 40 surgical procedures.

Surgical procedure and protocol

Oral hygiene instructions were provided, followed by scaling, polishing, and the elimination of identified causative factors. All surgical procedures were meticulously performed by a single, experienced surgeon (AG) using a standardized technique. Following the administration of 2% lidocaine local anesthesia containing 1:80,000 adrenaline, submarginal, cervical, and interdental incisions were meticulously executed to create a full-thickness flap. An apically repositioned flap was executed, ensuring a minimum of 2 mm of bone removal and the preservation of at least 2 mm of keratinized gingiva in




Figure 1. Flow diagram of the study

the same teeth bilaterally.¹¹ Adequate debridement was achieved through tissue resection and bone reduction. The surgical site was then meticulously cleaned and closed with Vicryl sutures (No. 04, Supa Medical Devices, Tehran, Iran).

Postoperatively, all the patients were prescribed 400 mg of ibuprofen daily (Jaber Ebne Hayan, Iran), a nonsteroidal anti-inflammatory medication. Additionally, 0.2% chlorhexidine was prescribed twice a day for one week. The patients were advised to consume only soft foods and to avoid any mechanical trauma to the treated areas for one week. Sutures were removed one week after the surgical procedure.

Interventions and laser irradiation

For both treatment groups, all the patients and the operator wore protective eyeglasses during the laser operation. In the case group, a 660 nm diode laser (Polaris 2, Astar Company, Bielsko-Biala, Poland) was applied continuously for 30 seconds to one side of the midline. This laser delivered 40 mW of power and an energy density of 1.2 J/cm² (Figure 2a). A diode laser (808 nm wavelength, 5 J/cm² energy, 200 mW power) was also applied continuously for 25 seconds (Figure 2b) in non-contact mode (Figure 3). Laser irradiation was performed twice on the soft tissue: immediately post-suture on the day of surgery and again three days later. The control side (the other side of the midline) underwent a placebo laser application using an identical technique and duration.

Outcomes

The primary objective of this study was to evaluate the impact of PBM on the early healing index (EHI) (Table 1) and postoperative pain following surgical crown lengthening. The EHI offers benefits, such as enabling the

assessment of initial repair within the first 24 hours after surgery. The EHI assesses three key factors: clinical signs of re-epithelialization (CSR), clinical signs of hemostasis (CSH), and clinical signs of inflammation (CSI).¹² This index was evaluated by two experienced periodontists (PB and RB) as the primary outcome measure on days 3 and 7 after surgery, with any discrepancies resolved through discussion.

Postoperative pain, a secondary outcome, was assessed by patients using a visual analog scale (VAS), which ranged from 0 (no pain) to 10 (unbearable pain).¹³ The researcher reminded patients to assess their pain levels on the day of surgery and again on days 1, 3, and 7 after surgery.

Randomization and concealment

In this study, individuals requiring crown lengthening surgery with at least two teeth in different quadrants were included. The assignment of intervention and control to each half-jaw was randomized using a closed opaque envelope method. For random allocation, two cards were labeled “left half-jaw intervention” and two others “right half-jaw intervention.” These four cards were placed into identical envelopes. For each patient, an assistant, responsible for laser application, opened one envelope to determine the treatment side.

Blinding

This study employed a triple-blind design, ensuring that the patient, outcome assessor, and data analyst remained unaware of the assignment to either the control or case groups.

Statistical analysis

Data underwent statistical analysis using SPSS 22. The




Figure 2. Laser instrument calibration. A. 660 nm beam. B. 808 nm beam




Figure 3. Laser application technique after crown lengthening surgery. A. 660 nm beam. B. 808 nm beam

Wilcoxon test was employed, with a two-sided significance level set at 0.05.

Results

Participants

This study evaluated 20 patients (mean age = 41.3 years; age range = 25–60 years). The cohort consisted of 11 males (73.3%) and 4 females (26.6%) (Figure 1). One patient was excluded from the study due to poor oral hygiene, and two patients did not attend their scheduled follow-up on day 3 after surgery.

Primary outcome

Table 2 presents the CSR grading for both groups on days 3 and 7 after surgery. No statistically significant differences were observed on either day 3 ($P=0.180$) or 7 ($P=1.0$) after surgery.

Table 3 illustrates the CSH grading on days 3 and 7 after surgery. A statistically significant difference was observed between the case and control groups on both day 3 ($P=0.03$) and day 7 ($P=0.01$).

Table 4 presents the CSI on days 3 and 7 after surgery. On day 3, no significant difference was observed in the

CSI grading between the two groups ($P=0.64$). However, by day 7, the case group demonstrated a significantly improved CSI grading ($P=0.004$).

Secondary outcome

Postoperative pain was significantly lower in the PBM group compared to the control group across all follow-up periods ($P<0.05$) (Table 5).

Discussion

The findings of this study suggest that a combined PBM protocol (a 660 nm laser at 40 mW, continuous, for 30 seconds, 1.2 J/cm² plus an 808 nm laser at 200 mW, continuous, for 25 seconds, 5 J/cm²) significantly expedited tissue healing and alleviated pain after surgical crown lengthening. Infrared radiation at 808 nm was used to induce analgesia, while red light at 660 nm was employed to facilitate tissue healing.¹⁴ Significant differences were observed between the two groups regarding CSI on day 7 and CSH on both days 3 and 7 ($P<0.05$). The case group consistently exhibited superior pain relief, as measured by the VAS score, at all follow-up time intervals ($P<0.05$).

Regarding soft tissue healing, the present study's

Table 1. The early healing index (EHI) scoring system

EHI subscales		Scores		
CSR	0: Visible distance between incision margins	3: Contact between incision margins	6: Merged incision margins	
CSH	0: Bleeding at the incision margins	1: Presence of fibrin on the incision margins	2: Absence of fibrin on the incision margins	
CSI	0: Redness involving >50% of the incision length and/or pronounced swelling	1: Redness 50% > incision length	2: Absence of redness along the incision length	

EHI: early healing index; CSR: clinical signs of re-epithelialization; CSH: clinical signs of hemostasis; CSI: clinical signs of inflammation

Table 2. Results of clinical signs of re-epithelialization (CSR)

Day	Group	Grade 0	Grade 3	Grade 6	P value
3 (n = 17)	Case	3 (18%)	14 (82%)	0	0.18
	Control	2 (12%)	13 (76%)	2 (12%)	
7 (n = 19)	Case	0	19 (100%)	0	1.0
	Control	0	19 (100%)	0	

Table 3. Results of clinical signs of hemostasis (CSH)

Day	Group	Grade 0	Grade 1	Grade 2	P value
3 (n = 17)	Case	0	8 (47%)	9 (53%)	0.03*
	Control	4 (24%)	8 (47%)	5 (29%)	
7 (n = 19)	Case	0	1	18	0.01*
	Control	0	7	12	

*Statistically significant.

Table 4. The results of clinical signs of inflammation (CSI)

Day	Group	Grade 0	Grade 1	Grade 2	P value
3 (n = 17)	Case	5 (29%)	7 (42%)	5 (29%)	0.64
	Control	6 (34%)	7 (42%)	4 (24%)	
7 (n = 19)	Case	1 (5%)	1 (5%)	17 (90%)	0.004*
	Control	0	10 (53%)	9 (47%)	

*Statistically significant.

Table 5. The results of postoperative pain (visual analog scale [VAS])

Day	Group	Mean	SD	P value
Day of surgery	Case	2.5	2.5	0.01*
	Control	3.4	2.9	
1	Case	1.5	1.9	0.005*
	Control	2.7	2.9	
3	Case	0.7	1.2	0.004*
	Control	1.8	2.2	
7	Case	0.3	0.8	0.03*
	Control	0.6	1.0	

SD: standard deviation; * Statistically significant.

findings align with previous research by Amorim et al³ (685 nm, 50 mW, 4 J/cm), Kohale et al¹⁵ (940 nm, 100 mW), Pejic et al¹⁶ (670 nm), Lingamaneni et al⁶ (810 nm, 0.1 W, 5 min), Ozturan et al⁸ (588 nm, 128 mW, 5 min, 4 J), and Ustaoglu et al¹⁷ (940 nm).

Amorim et al³ previously investigated the use of LLLT following gingivectomy, concluding that laser application enhanced clinical repair. However, their findings may be subject to bias due to the use of dressings, which can independently influence healing and pain relief. Similarly, Kohale et al¹⁵ reported the effectiveness of PBM in promoting healing after gingivectomy. A significant number of clinical samples were included in this study, and similar to the current investigation, no dressing was used. Pejic et al¹⁶ investigated the impact of PBM following treatment for chronic mild periodontitis, concluding that laser application enhanced both clinical symptoms and healing outcomes. Similarly, Lingamaneni et al⁶ observed improved gingival epithelialization after gingivectomy. A limitation of their study was the restricted sample size and the specific postoperative dressing employed. Despite the small sample size, Ozturan et al⁸ similarly reported that laser application accelerated repair in coronally advanced flaps. Furthermore, Ustaoglu et al¹⁷ concluded that LLLT improved wound healing at the donor site of free gingival grafts and helped preserve tissue thickness in those areas.

The capacity of PBM to influence inflammation and enhance healing likely stems from its impact on the initial phases of wound healing. The early postoperative period is crucial for wound healing, as inflammatory cells play a vital role in this stage, clearing tissue debris and facilitating the migration of keratinocytes and fibroblasts. Postoperative recovery relies on various gingival cells, including fibroblasts, keratinocytes, and immune cells. The healing cascade involves a series of events orchestrated by cytokines and growth factors released by immune cells.

The findings of the repair process in the current study diverge from those reported by Damante et al,¹⁸ Ozcelik et al,⁷ and Ravi et al.¹⁹ This inconsistency can be attributed to methodological differences, specifically variations in laser protocols, surgical techniques, and the limited number of participants in these studies.

Damante et al¹⁸ observed no positive outcomes in evaluating the impact of a 670 nm Ga-Al-As laser on

tissue repair following gingivectomy. This lack of efficacy may be attributed to several factors, including the use of a power output of <15 mW. Research suggests that greater power and wavelengths within the red spectrum are necessary to accelerate tissue repair.¹⁴ Ozcelik et al⁷ investigated the impact of PBM following a gingivectomy, observing no significant intergroup differences in tissue repair. This discrepancy may be attributable to variations in the surgical procedures themselves and the nature of secondary repair after gingivectomy. Surgical wounds inherently differ considerably in terms of the type of surgery, wound depth, and the subsequent recovery protocol.

PBM is recommended to alleviate patient discomfort and complaints stemming from postoperative pain. The pain-relieving effects observed after laser application may be attributed to the accelerated wound healing process. This acceleration can be explained by enhanced keratinocyte migration, expedited epithelialization, and increased fibroblast proliferation and neovascularization.^{20,21}

Regarding pain outcomes, the findings of this study align with those reported by Doshi et al,²⁰ Lafzi et al,²¹ Etemadi et al,²² Ravi et al,¹⁹ Heidari et al,²³ and Sadighi et al.²⁴ Similarly, Madi and Mahmoud²⁵ investigated the impact of a 660 nm diode laser following gingivectomy on 20 patients, demonstrating improved repair and reduced pain within the laser-treated group. This study's methodology introduced potential confounding factors due to the use of dressings that could influence healing outcomes, as well as the application of foil beneath the dressing, which may stimulate the surgical site. In contrast, Almeida et al⁹ investigated the impact of PBM on 10 patients following FG and concluded that laser therapy did not effectively reduce pain or accelerate healing. Heidari et al⁵ investigated the effect of laser therapy on FG repair and associated pain. Their findings indicated accelerated healing in the case group, but reported similar pain levels between both groups. This contrasts with the results of the current study, a discrepancy that could be attributed to the smaller sample size and differing surgical techniques employed. PBM, particularly within the energy range of 4–20 J/cm², shows promise in alleviating pain after periodontal surgery.²⁶ However, further clinical trials employing similar parameters are essential to establish the optimal dose and clinical protocol.

Research investigating the application of PBM in crown lengthening surgery is limited. Consequently, more rigorously designed studies, featuring larger sample sizes and diverse clinical parameters, are essential to draw comprehensive conclusions regarding its efficacy. Furthermore, the specific type of surgical procedure may significantly influence the effectiveness of lasers in mitigating pain and promoting accelerated healing.

Conclusion

PBM has been shown to enhance the healing process and reduce pain following crown lengthening surgery.

Specifically, PBM significantly improved the CSI by day 7 and the CSH by days 3 and 7 after surgery. Furthermore, it effectively alleviated pain on the day of surgery, as well as on days 1, 3, and 7 after surgery.

Acknowledgments

We gratefully acknowledge the continuous guidance and support provided by the Vice-Chancellor for Research at Isfahan University of Medical Sciences.

Authors' Contribution

Conceptualization: Parichehr Behfarnia.

Data curation: Parichehr Behfarnia, Reza Birang.

Formal analysis: Amineh Ghaznavi, Seyed Amir Mirghaderi .

Funding acquisition: Parichehr Behfarnia.

Investigation: All authors.

Methodology: Parichehr Behfarnia, Reza Birang.

Project administration: All authors.

Resources: Amineh Ghaznavi, Seyed Amir Mirghaderi.

Supervision: Parichehr Behfarnia, Reza Birang.

Validation: Parichehr Behfarnia.

Visualization: Amineh Ghaznavi, Seyed Amir Mirghaderi.

Writing-original draft: Parichehr Behfarnia, Amineh Ghaznavi, Seyed Amir Mirghaderi.

Writing-review & editing: Parichehr Behfarnia, Seyed Amir Mirghaderi.

Competing Interests

The authors declare that they have no conflicts of interest.

Consent for Publication

Not applicable.

Data Availability Statement

The datasets generated during the current study are available from the corresponding author on reasonable request.

Ethical Approval

The present study was approved by the Ethics Committee of Isfahan University of Medical Sciences (IR.MUI.RESEARCH.REC.1400.128). All procedures performed in studies involving human participants were in accordance with the ethical standards of the institutional and/or national research committee and with the 1964 Helsinki Declaration and its later amendments, or with comparable ethical standards. This study was registered at the Iranian Registry of Clinical Trials (IRCT) with registration number IRCT20110109005570N12.

Funding

This study was funded by the Vice-Chancellor for Research at Isfahan University of Medical Sciences (Grant No. 530080).

References

- Blyleven GM, Johnson TM, Inouye KA, Stancoven BW, Lincicum AR. Factors influencing intraoperative and postoperative complication occurrence: a series of 1135 periodontal and implant-related surgeries. *Clin Exp Dent Res*. 2024;10(1):e849. doi: [10.1002/cre2.849](#).
- Pippi R. Post-surgical clinical monitoring of soft tissue wound healing in periodontal and implant surgery. *Int J Med Sci*. 2017;14(8):721-8. doi: [10.7150/ijms.19727](#).
- Amorim JC, de Sousa GR, de Barros Silveira L, Prates RA, Pinotti M, Ribeiro MS. Clinical study of the gingiva healing after gingivectomy and low-level laser therapy. *Photomed Laser Surg*. 2006;24(5):588-94. doi: [10.1089/pho.2006.24.588](#).
- Bahrani H, Moharrami A, Mirghaderi P, Mortazavi SM. Low-level laser and light therapy after total knee arthroplasty improves postoperative pain and functional outcomes: a three-arm randomized clinical trial. *Arthroplast Today*. 2023;19:101066. doi: [10.1016/j.artd.2022.10.016](#).
- Heidari M, Paknejad M, Jamali R, Nokhbatolfighahaei H, Fekrazad R, Moslemi N. Effect of laser photobiomodulation on wound healing and postoperative pain following free gingival graft: a split-mouth triple-blind randomized controlled clinical trial. *J Photochem Photobiol B*. 2017;172:109-14. doi: [10.1016/j.jphotobiol.2017.05.022](#).
- Lingamaneni S, Mandadi LR, Pathakota KR. Assessment of healing following low-level laser irradiation after gingivectomy operations using a novel soft tissue healing index: a randomized, double-blind, split-mouth clinical pilot study. *J Indian Soc Periodontol*. 2019;23(1):53-7. doi: [10.4103/jisp.jisp_226_18](#).
- Ozcelik O, Cenk Haytac M, Kunin A, Seydaoglu G. Improved wound healing by low-level laser irradiation after gingivectomy operations: a controlled clinical pilot study. *J Clin Periodontol*. 2008;35(3):250-4. doi: [10.1111/j.1600-051X.2007.01194.x](#).
- Ozturan S, Durukan SA, Ozcelik O, Seydaoglu G, Cenk Haytac M. Coronally advanced flap adjunct with low intensity laser therapy: a randomized controlled clinical pilot study. *J Clin Periodontol*. 2011;38(11):1055-62. doi: [10.1111/j.1600-051X.2011.01774.x](#).
- Almeida AL, Esper LA, Sbrana MC, Ribeiro IW, Kaizer RO. Utilization of low-intensity laser during healing of free gingival grafts. *Photomed Laser Surg*. 2009;27(4):561-4. doi: [10.1089/pho.2008.2292](#).
- Schulz KF, Altman DG, Moher D. CONSORT 2010 statement: updated guidelines for reporting parallel group randomized trials. *Ann Intern Med*. 2010;152(11):726-32. doi: [10.7326/0003-4819-152-11-201006010-00232](#).
- Cairo F, Graziani F, Franchi L, Defraia E, Pini Prato GP. Periodontal plastic surgery to improve aesthetics in patients with altered passive eruption/gummy smile: a case series study. *Int J Dent*. 2012;2012:837658. doi: [10.1155/2012/837658](#).
- Marini L, Rojas MA, Sahrman P, Aghazada R, Pilloni A. Early Wound Healing Score: a system to evaluate the early healing of periodontal soft tissue wounds. *J Periodontal Implant Sci*. 2018;48(5):274-83. doi: [10.5051/jpis.2018.48.5.274](#).
- Yen CA, Griffin TJ, Cheung WS, Chen J. Effects of platelet concentrate on palatal wound healing after connective tissue graft harvesting. *J Periodontol*. 2007;78(4):601-10. doi: [10.1902/jop.2007.060275](#).
- Gholami L, Asefi S, Hooshyarfarad A, Sculean A, Romanos GE, Aoki A, et al. Photobiomodulation in periodontology and implant dentistry: part 2. *Photobiomodul Photomed Laser Surg*. 2019;37(12):766-83. doi: [10.1089/photob.2019.4731](#).
- Kohale BR, Agrawal AA, Raut CP. Effect of low-level laser therapy on wound healing and patients' response after scalpel gingivectomy: a randomized clinical split-mouth study. *J Indian Soc Periodontol*. 2018;22(5):419-26. doi: [10.4103/jisp.jisp_239_18](#).
- Pejicic A, Kojovic D, Kesic L, Obradovic R. The effects of low-level laser irradiation on gingival inflammation. *Photomed Laser Surg*. 2010;28(1):69-74. doi: [10.1089/pho.2008.2301](#).
- Ustaoglu G, Ercan E, Tunali M. Low-level laser therapy in enhancing wound healing and preserving tissue thickness at free gingival graft donor sites: a randomized, controlled clinical study. *Photomed Laser Surg*. 2017;35(4):223-30. doi: [10.1089/pho.2016.4163](#).
- Damante CA, Greggi SL, Sant'ana AC, Passanezi E. Clinical evaluation of the effects of low-intensity laser (GaAlAs) on wound healing after gingivoplasty in humans. *J Appl Oral Sci*. 2004;12(2):133-6. doi: [10.1590/s1678-77572004000200010](#).
- Ravi M, Nalini H, Prasad A, Devi R. The effect of biostimulation with 980-nm diode laser on postoperative pain and tissue

- response after surgical periodontal therapy: a pilot study. *J Indian Acad Dent Spec Res*. 2016;3(2):47. doi: [10.4103/jiadsr.jiadsr_9_17](https://doi.org/10.4103/jiadsr.jiadsr_9_17).
20. Doshi S, Jain S, Hegde R. Effect of low-level laser therapy in reducing dentinal hypersensitivity and pain following periodontal flap surgery. *Photomed Laser Surg*. 2014;32(12):700-6. doi: [10.1089/pho.2014.3802](https://doi.org/10.1089/pho.2014.3802).
21. Lafzi A, Kadkhodazadeh M, Mojahedi SM, Amid R, Shidfar S, Baghani MT. The clinical evaluation of the effects of low-level laser therapy on the donor and recipient sites of the free gingival graft: a case series. *J Lasers Med Sci*. 2019;10(4):355-60. doi: [10.15171/jlms.2019.58](https://doi.org/10.15171/jlms.2019.58).
22. Etemadi A, Sadeghi M, Dadjou MH. The effects of low-level 660-nm laser irradiation on pain and teeth hypersensitivity after periodontal surgery. *J Lasers Med Sci*. 2011;2(3):103-8.
23. Heidari M, Fekrazad R, Sobouti F, Moharrami M, Azizi S, Nokhbatolfoghahaei H, et al. Evaluating the effect of photobiomodulation with a 940-nm diode laser on post-operative pain in periodontal flap surgery. *Lasers Med Sci*. 2018;33(8):1639-45. doi: [10.1007/s10103-018-2492-y](https://doi.org/10.1007/s10103-018-2492-y).
24. Sadighi M, Faramarzi M, Chitsazha R, Ghasemi Rad M, Ranjbar S. Effect of photobiomodulation on pain control after clinical crown lengthening surgery. *J Adv Periodontol Implant Dent*. 2021;13(2):91-4. doi: [10.34172/japid.2021.014](https://doi.org/10.34172/japid.2021.014).
25. Madi M, Mahmoud MM. The evaluation of healing effect of low-level laser treatment following gingivectomy. *Beni Suef Univ J Basic Appl Sci*. 2020;9(1):25. doi: [10.1186/s43088-020-00053-z](https://doi.org/10.1186/s43088-020-00053-z).
26. Wagner VP, Meurer L, Martins MA, Danilevicz CK, Magnusson AS, Marques MM, et al. Influence of different energy densities of laser phototherapy on oral wound healing. *J Biomed Opt*. 2013;18(12):128002. doi: [10.1117/1.Jbo.18.12.128002](https://doi.org/10.1117/1.Jbo.18.12.128002).